


Short answer questions (8 points each):

1. Express the triple integral of the function $x^2 - y + z$ over the region bounded by the planes $x = 0, y = 0, z = 0$, and $2x + y + z = 8$. Do not evaluate the iterated integral.

$$z = 8 - 2x - y$$

$$\int_0^4 \int_0^{8-2x} \int_0^{8-2x-y} x^2 - y + z \, dz \, dy \, dx$$



2. Is the vector field $\vec{F}(x, y, z) = \langle xy, xz, yz \rangle$ conservative? If so, find a potential function.
If not, demonstrating that it is not completely answers the question


Test by finding curl:

$$\nabla \times \vec{F} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ xy & xz & yz \end{vmatrix} = (z-x)\vec{i} - (0-0)\vec{j} + (z-x)\vec{k} \neq \vec{0}$$

so it's not conservative

Other orders of integration for #1


$$\int_0^8 \int_0^{4-y/2} \int_0^{8-2x-y} x^2 - y + z \, dz \, dx \, dy$$



$$\int_0^4 \int_0^{8-2x} \int_0^{8-2x-2} x^2 - y + z \, dy \, dz \, dx$$

balk;
 $y = 8 - 2x - z$


$$\int_0^8 \int_0^{4-y/2} \int_0^{8-2x-2} x^2 - y + z \, dy \, dx \, dz$$



$$\int_0^8 \int_0^{8-y} \int_{4-y/2-z/2}^{4-y/2} x^2 - y + z \, dx \, dz \, dy$$

right side;
 $x = 4 - y/2 - z/2$

$$\int_0^8 \int_0^{8-y} \int_0^{4-y/2-z/2} x^2 - y + z \, dx \, dy \, dz$$



Short answer questions, continued.

3. Let C be the portion of the circle of radius 1 about the origin lying in the first quadrant beginning at $(0, 1)$ and ending at $(1, 0)$. Give a parametrization of C and use it to express the line integral

$$\int_C \vec{F} \cdot d\vec{r}$$

as an ordinary definite integral, where \vec{F} is the vector field $\vec{F}(x, y) = \langle -1, x \rangle$. Do not evaluate the resulting definite integral.


parametrization by γ

$$\vec{\gamma}(t) = \langle \cos t, \sin t \rangle \quad t \in [0, \pi/2]$$

$$\vec{\gamma}'(t) = \langle -\sin t, \cos t \rangle$$

$$\vec{F}(\vec{\gamma}(t)) = \langle -1, \cos t \rangle$$

$$\begin{aligned} \text{So } \int_C \vec{F} \cdot d\vec{r} &= \int_0^{\pi/2} (-1)(-\sin t) + (\cos t)(\cos t) dt \\ &= \int_0^{\pi/2} \sin t + \cos^2 t dt \end{aligned}$$



4. Express the volume of the region described in the next sentence as an iterated integral by using cylindrical coordinates. Do not evaluate the iterated integral.


The region lies between the planes $z = 1$ and $z = -1$, inside the sphere of radius 4 about the origin and outside the cylinder of radius 2 about the z -axis.

$$2 \leq r \leq \sqrt{16 - z^2}$$

$$-1 \leq z \leq 1$$

$$0 \leq \theta \leq 2\pi$$

$$dV = r dr dz d\theta$$



$$V = \iiint_R dV = \int_0^{2\pi} \int_{-1}^1 \int_2^4 r dr dz d\theta$$

#3 with parametrization by $t = y$

$$\vec{r}(t) = \left\langle \sqrt{1-t^2}, t \right\rangle \quad t \in [0, 1]$$

$$\begin{aligned}\vec{r}'(t) &= \left\langle \frac{1}{2} (1-t^2)^{-\frac{1}{2}} \cdot (-2t), 1 \right\rangle \\ &= \left\langle \frac{-t}{\sqrt{1-t^2}}, 1 \right\rangle\end{aligned}$$

$$\vec{F}(\vec{r}(t)) = \left\langle -1, \sqrt{1-t^2} \right\rangle$$

So

$$\int_C \vec{F} \cdot d\vec{r} = \int_0^1 (-1) \frac{-t}{\sqrt{1-t^2}} + 1 \sqrt{1-t^2} dt$$

$$= \int_0^1 \frac{t}{\sqrt{1-t^2}} + \sqrt{1-t^2} dt$$

Yet more short answer questions.

5. Find the divergence of $\vec{Q}(x, y, z) = ye^z\vec{i} + ze^x\vec{j} + xe^y\vec{k}$, and use it to decide whether or not the vector field could represent the velocity of a flow in an incompressible fluid.

$$\nabla \cdot \vec{Q} = \frac{\partial}{\partial x} ye^z + \frac{\partial}{\partial y} ze^x + \frac{\partial}{\partial z} xe^y$$
$$= 0 + 0 + 0 = 0$$

Yes it could represent velocities in
an incompressible fluid flow

6. Find the curl of the vector field $\vec{Q}(x, y, z) = ye^z\vec{i} + ze^x\vec{j} + xe^y\vec{k}$, and use it to decide whether the vector field is irrotational.

$$\nabla \times \vec{Q} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ ye^z & ze^x & xe^y \end{vmatrix}$$
$$= (xe^y - e^x)\vec{i} - (e^y - ye^z)\vec{j} + (ze^x - e^x)\vec{k}$$

No it is not irrotational.

Long questions (18 points each)

7. Let R be the region lying in the first octant (all coordinates positive) and inside the sphere of radius 3 about the origin. Use spherical coordinates to express the following triple integral as an iterated integral and evaluate the resulting iterated integral.


$$\iiint_R \frac{z}{x^2 + y^2 + z^2} dV$$

$$x^2 + y^2 + z^2 = \rho^2$$

$$z = \rho \cos \varphi$$

$$dV = \rho^2 \sin \varphi \, d\rho \, d\varphi \, d\theta$$

$$\int_0^{\pi/2} \int_0^{\pi/2} \int_0^3 \frac{\rho \cos \varphi \, \rho^2 \sin \varphi}{\rho^2} \, d\rho \, d\varphi \, d\theta$$



$$= \int_0^{\pi/2} \int_0^{\pi/2} \int_0^3 \rho \cos \varphi \sin \varphi \, d\rho \, d\varphi \, d\theta$$

$$= \int_0^{\pi/2} d\theta \int_0^{\pi/2} \cos \varphi \sin \varphi \, d\varphi \int_0^3 \rho \, d\rho$$

Since it's a ρ, φ, θ parameter
space of a $1/8$ th solid
and the integrand factors
as $1/r^2$ function

$$\begin{aligned} & \text{let } u = \sin \varphi \\ & du = \cos \varphi \, d\varphi \\ & \varphi = 0 \Rightarrow u = 0 \quad \varphi = \pi/2 \Rightarrow u = 1 \end{aligned}$$

$$= \frac{\pi}{2} \int_0^1 u \, du \cdot \left[\frac{\rho^2}{2} \right]_0^3$$

$$= \frac{\pi}{2} \cdot \frac{u^2}{2} \Big|_0^1 \cdot \left[\frac{\rho^2}{2} \right]_0^3 = \frac{\pi}{2} \cdot \frac{1}{2} \cdot \frac{9}{2} = \frac{9\pi}{8}$$

8. Find the average value of the function $w(x, y) = x^2$ on the disk of radius 1 centered at the origin (in the xy -plane).

$$\text{Area} = \pi \cdot 1^2 = \pi$$

$$\text{so Avg value} = \frac{1}{\pi} \iint_R x^2 dA$$

polar coords seems good for this: since region is
 $0 \leq \theta \leq 2\pi$
 $0 \leq r \leq 1$

$$\hookrightarrow x^2 = r^2 \cos^2 \theta$$

$$dA = r dr d\theta$$

$$\text{Avg} = \frac{1}{\pi} \int_0^{2\pi} \int_0^1 r^3 \cos^2 \theta dr d\theta$$

$$= \frac{1}{\pi} \int_0^{2\pi} \cos^2 \theta d\theta \int_0^1 r^3 dr$$

$$= \frac{1}{\pi} \left[\frac{\theta}{2} + \frac{\sin 2\theta}{4} \right] \Big|_0^{2\pi} \cdot \frac{r^4}{4} \Big|_0^1$$

using #14
in the
integral table

$$= \frac{1}{\pi} \left[\frac{2\pi}{2} + 0 - (0-0) \right] \cdot \left(\frac{1}{4} - 0 \right)$$

$$= \frac{1}{\pi} \cdot \pi \cdot \frac{1}{4} = \frac{1}{4}$$

#8 In rectangular coords : region is

$$-\sqrt{1-x^2} \leq y \leq \sqrt{1-x^2}$$

$$-1 \leq x \leq 1$$

So

$$\text{Avg} = \frac{1}{\pi} \int_{-1}^1 \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} x^2 dy dx$$

$$= \frac{1}{\pi} \int_{-1}^1 x^2 y \Big|_{y=-\sqrt{1-x^2}}^{y=\sqrt{1-x^2}} dx$$

$$= \frac{1}{\pi} \int_{-1}^1 2x^2 \sqrt{1-x^2} dx = \frac{2}{\pi} \int_{-1}^1 x^2 \sqrt{1-x^2} dx \quad \begin{array}{l} \text{by form} \\ \text{for} \\ \text{integ'n} \\ \text{table} \end{array}$$

$$= \frac{2}{\pi} \left[-\frac{x}{4} (1-x^2)^{3/2} + \frac{1}{8} (x\sqrt{1-x^2} + \arcsin x) \right]_{x=-1}^1$$

doubled since it's an odd function

$$= \frac{4}{\pi} \left[-\frac{1}{4} \cdot 0 + \frac{1}{8} \cdot (1 \cdot 0 + \frac{\pi}{2}) \right]$$

and
we're subtracting
its value at -1

from its value at 1

$$= \frac{4}{\pi} \cdot \frac{1}{8} \cdot \frac{\pi}{2} = \frac{1}{4}$$

evaluated
at $x=1$

9. Find the average value of the function $w(x, y) = x^2$ on the circle of radius 1 centered at the origin (in the xy -plane). (Just on the circle, not the whole disk – that was the previous question.)

length of curve \Rightarrow circumference $2\pi \cdot 1 = 2\pi$

$$\text{so avg value} = \frac{1}{2\pi} \int_C x^2 ds$$

parametrize by the circle

$$\vec{r}(t) = \langle \cos t, \sin t \rangle \quad t \in [0, 2\pi]$$

$$\text{so } \vec{r}'(t) = \langle -\sin t, \cos t \rangle$$

$$|\vec{r}'(t)| = \sqrt{(-\sin t)^2 + (\cos t)^2} = 1$$

$$\text{so } ds = dt$$

$$x^2 = \cos^2 t$$

$$\therefore \text{Avg value} = \frac{1}{2\pi} \int_0^{2\pi} \cos^2 t dt$$

$$= \frac{1}{2\pi} \left[\frac{t}{2} + \frac{\sin 2t}{4} \right] \Big|_0^{2\pi} \quad \begin{matrix} \hookrightarrow \\ \text{from 17a,} \\ \text{stay-fbk.} \end{matrix}$$

$$= \frac{1}{2\pi} \left[\frac{2\pi}{2} + 0 - (0+0) \right]$$

$$= \frac{1}{2}$$

10. A large magnetic sheet (idealized as) occupying the plane $z = 4$ (in a convenient coordinate system for the problem with distances in meters) exerts a force of $\frac{1}{(4-z)^2} \vec{k}$ Newtons on an iron particle located at (x, y, z) when $z < 4$. If the force causes the iron particle, constrained in a helical tube, to move from $(3, 0, 0)$ to $(3, 0, 2)$ along the curve given parametrically by $\vec{r}(t) = \langle 3 \cos t, 3 \sin t, t/\pi \rangle$. Find the work done by the force in Nm.

If will be easier if it's conservative. Let's test:

$$\text{If } \vec{F} = \frac{1}{(4-z)^2} \vec{k}$$

$$\nabla \times \vec{F} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 0 & 0 & (4-z)^{-2} \end{vmatrix} = (0-0)\vec{i} - (0-0)\vec{j} + (0-0)\vec{k}$$

Since it's conservative find a potential function!

$$f = \int (4-z)^{-2} dz = + (4-z)^{-1} + C(x, y)$$

plausibly, $f = + (4-z)^{-1}$ works since the x and y partials are 0

$$\begin{aligned} \text{So } W &= \int_C \vec{F} \cdot d\vec{r} = f(3, 0, 2) - f(3, 0, 0) \\ &= + (4-2)^{-1} - + (4-0)^{-1} \\ &= + \frac{1}{2} - \frac{1}{4} \\ &= - \frac{1}{4} \text{ Nm} \end{aligned}$$

#10 by finding the line integral directly

curve $\vec{r}(t) = \langle 3\cos t, 3\sin t, t/\pi \rangle \quad t \in [0, 2\pi]$

$$\vec{r}'(t) = \langle -3\sin t, 3\cos t, 1/\pi \rangle$$

$$F(\vec{r}(t)) = \frac{1}{(4 - t/\pi)^2} \vec{k}$$

$$S_0 \int_0^{2\pi} \langle 0, 0, (4 - t/\pi)^{-2} \rangle \cdot \langle -3\sin t, 3\cos t, 1/\pi \rangle dt$$

$$\begin{aligned} &= \int_0^{2\pi} \frac{1}{\pi} (4 - t/\pi)^{-2} dt \quad \text{let } u = 4 - t/\pi \\ &\qquad \qquad \qquad du = -\frac{1}{\pi} dt \\ &\qquad \qquad \qquad \therefore -du = \frac{1}{\pi} dt \\ &= \int_4^2 -u^{-2} du \quad t \geq 0 \Rightarrow u = 4 \\ &\qquad \qquad \qquad t = 2\pi \Rightarrow u = 4 - \frac{2\pi}{\pi} = 2 \end{aligned}$$

$$= \int_2^4 u^{-2} du = -u^{-1} \Big|_2^4$$

$$= -\frac{1}{4} - (-\frac{1}{2}) = \frac{1}{4} \text{ Nm}$$