

NAME _____

Rec. Instructor: _____

Signature _____

Rec. Time _____

CALCULUS III - PRACTICE TEST FINAL

Show all work for full credit. No books or notes are permitted.

Problem	Points	Possible	Problem	Points	Possible
1			7		
2			8		
3			9		
4			10		
5			11		
6			EC		
Total Score					

Note: Bold letters, like **u**, are considered vectors unless specified otherwise.

You are free to use the following formulas on any of the problems.

Projection: $\text{proj}_{\mathbf{u}} \mathbf{v} = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\|^2} \mathbf{u}$

Cylindrical Coordinates:

$$x = r \cos(\theta)$$

$$y = r \sin(\theta)$$

$$z = z$$

$$dV = r dr d\theta dz$$

$$r = \sqrt{x^2 + y^2}$$

$$\tan(\theta) = \frac{y}{x}$$

$$z = z$$

Spherical Coordinates:

$$x = \rho \cos(\theta) \sin(\varphi)$$

$$y = \rho \sin(\theta) \sin(\varphi)$$

$$z = \rho \cos(\varphi)$$

$$dV = \rho^2 \sin(\phi) d\rho d\theta d\phi$$

$$\rho = \sqrt{x^2 + y^2 + z^2}$$

$$\tan(\theta) = \frac{y}{x}$$

$$\cos(\varphi) = \frac{z}{\rho}$$

Second Derivative Test: Let $z = f(x, y)$ be a function of two variables for which the first- and second-order partial derivatives are continuous on some disk containing the point (x_0, y_0) . Suppose $f_x(x_0, y_0) = 0$ and $f_y(x_0, y_0) = 0$. Define the quantity

$$D = f_{xx}(x_0, y_0)f_{yy}(x_0, y_0) - (f_{xy}(x_0, y_0))^2$$

- i. If $D > 0$ and $f_{xx}(x_0, y_0) > 0$, then f has a local minimum at (x_0, y_0) .
- ii. If $D > 0$ and $f_{xx}(x_0, y_0) < 0$, then f has a local maximum at (x_0, y_0) .
- iii. If $D < 0$, then f has a saddle point at (x_0, y_0) .
- iv. If $D = 0$, then the test is inconclusive.

Trig Identities: $\cos^2(x) = \frac{1}{2} + \frac{1}{2} \cos(2x) \quad \sin^2(x) = \frac{1}{2} - \frac{1}{2} \cos(2x)$

Line Integrals:

$$\int_C f ds = \int_a^b f(\mathbf{r}(t)) \|\mathbf{r}'(t)\| dt$$


$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_a^b \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt$$

Surface Integrals:

$$\int \int_S f dS = \int \int_R f(\mathbf{r}(u, v)) \|(\mathbf{t}_u \times \mathbf{t}_v)\| du dv$$

$$\int \int_S \mathbf{F} \cdot d\mathbf{S} = \int \int_R \mathbf{F}(\mathbf{r}(u, v)) \cdot (\mathbf{t}_u \times \mathbf{t}_v) du dv$$

1. For this problem we refer to the following diagram, which is drawn to scale:



The vectors \mathbf{A} , \mathbf{B} , \mathbf{C} , \mathbf{D} , \mathbf{E} , \mathbf{F} , and \mathbf{G} all have length three. All of the angles between the vectors are multiples of 30 degrees. Compute the following explicitly:

- a) $\mathbf{A} \cdot \mathbf{E}$
- b) $\|\mathbf{B} \times \mathbf{F}\|$
- c) $\mathbf{C} \cdot \mathbf{E}$
- d) $\|\mathbf{D} - \mathbf{G}\|$
- e) $\mathbf{A} \cdot \mathbf{A}$

- 2.** Use Lagrange Multipliers to find the maximum and minimum of the following function

$$f(x, y, z) = x^2 + 3z^2$$

subject to the constraint

$$x^2 + y^2 + 4z^2 = 36.$$

- 3.** Let E be the region given by $[0, 1] \times [0, 2] \times [0, 3]$. Compute the triple integral

$$\int \int \int_E z + xe^{xy} dV.$$

4. Calculate the following line integrals:

a) $\int_C f ds$ where $f(x, y, z) = 3x - yz$ and C is the line between the points $(1, 3, 0)$ and $(2, 5, 4)$.

b) $\int_C \mathbf{F} \cdot d\mathbf{r}$ where $\mathbf{F}(x, y) = \langle 6y, 2 \rangle$ and C is the piece of the graph $y = x^2$ between $x = 0$ and $x = 3$.

5. Calculate $\int_C \mathbf{F} \cdot d\mathbf{r}$ where $\mathbf{F}(x, y) = (y^2 - 4y + 5)\mathbf{i} + (2xy - 4x + 9)\mathbf{j}$ on the upper half of the ellipse $\frac{x^2}{9} + \frac{y^2}{25} = 1$, oriented clockwise. State any theorems used.

- 6.** Let E be the region such that $\sqrt{x^2 + y^2} \leq z \leq 3$ and $x \geq 0$. Consider

$$\int \int \int_E x dV.$$

- a)** If you were evaluating the given integral over E , would you integrate in Cartesian, Polar, Cylindrical, or Spherical Coordinates? Explain why you choose the coordinate system you did, and express the bounds for the region E in that system.

- b)** Evaluate

$$\int \int \int_E x dV.$$

7. Calculate the line integral

$$\oint_C (6y + x^2)dx + (1 - 2xy)dy$$

where C is the boundary of the upper half of the unit circle (this includes the x -axis), oriented counterclockwise. State any theorems used.

8. Evaluate the integral

$$\iint_S xz dS$$

where S is the portion of the sphere of radius 1 where $x \leq 0$, $y \geq 0$, $z \leq 0$.

9. Compute the integral

$$\iint_S \operatorname{curl} \mathbf{F} \cdot d\mathbf{S}$$

where $\mathbf{F}(x, y, z) = xz\mathbf{i} + yz\mathbf{j} + xz\mathbf{k}$ and S is the part of the sphere $x^2 + y^2 + z^2 = 9$ that lies in the cylinder $x^2 + y^2 = 1$ and above the xy -plane. State any theorems used.

10. For the following, determine if the vector field is conservative or not. If it is, find a potential function.

a) $\mathbf{F}(x, y, z) = \langle \frac{2}{3}y^3z^2, 2xy^2z^2, x^2y^2z^2 \rangle.$

b) $\mathbf{F}(x, y) = \langle 6x^2 - 2xy^2 + \frac{y}{2\sqrt{x}}, -2x^2y + 4 + \sqrt{x} \rangle.$

11. Answer the following short answer questions.

a) Suppose that $\nabla f(1, 1) = 0$, and $f_{xx}(1, 1)$ and $f_{yy}(1, 1)$ are both positive. Do you need to know any more information to determine if $(1, 1)$ is a local minimum?

b) Suppose that $\nabla g(0, 0) = 0$, and the discriminant D of g is 0 at $(0, 0)$. What does the Second Derivative Test tell us about how g behaves at $(0, 0)$?

c) If I set up an integral that looks like:

$$\int_0^z \int_2^{-1} \int_0^{y+3} f(x, y, z) dz dx dy$$

what is the problem with my integral set up?

d) Find the Jacobian of the transformation $x(\rho, \theta, \varphi) = \rho \cos(\theta) \sin(\varphi)$, $y(\rho, \theta, \varphi) = \rho \sin(\theta) \sin(\varphi)$, and $z(\rho, \theta, \varphi) = \rho \cos(\varphi)$.

e) Find the equation of the tangent plane to $f(x, y) = y \cos(x) + \sin(xy)$ at point $(\frac{\pi}{4}, 2)$.

f) If f is a function of x , y , z , and x , y , and z are each functions of u , v and w , use the chain rule to express $\frac{\partial f}{\partial u}$ and $\frac{\partial f}{\partial w}$.

g) Compute the divergence and curl of $\mathbf{F} = \langle x^2y, xyz, -x^2z^2 \rangle$.

h) Calculate the limit if it exists. If the limit does not exist, explain why not.

$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy^2}{x^2 + 3y^4}$$

EC. Use the Divergence Theorem to evaluate $\int \int_S \mathbf{F} \cdot d\mathbf{S}$ where

$$\mathbf{F}(x, y, z) = \langle x^2, y + z, xy \rangle$$

and S is the sphere of radius 3 with $z \leq 0$ and $y \leq 0$. Note that all three surfaces of this solid are included in S .