

NAME \_\_\_\_\_

Rec. Instructor: \_\_\_\_\_

Signature \_\_\_\_\_

Rec. Time \_\_\_\_\_

## CALCULUS III - EXAM 3

Show all work for full credit. No books or notes are permitted.

Problem	Points	Possible
1		15
2		20
3		10
4		20
5		20
6		15
Total Score		100

Note: Bold letters, like  $\mathbf{u}$ , are considered vectors unless specified otherwise.

You are free to use the following formulas on any of the problems.

**Cylindrical Coordinates:**

$$x = r \cos(\theta)$$

$$r = \sqrt{x^2 + y^2}$$

$$y = r \sin(\theta)$$

$$\tan(\theta) = \frac{y}{x}$$

$$z = z$$

$$z = z$$

$$dV = r dr d\theta dz$$

**Spherical Coordinates:**

$$x = \rho \cos(\theta) \sin(\varphi)$$

$$\rho = \sqrt{x^2 + y^2 + z^2}$$

$$y = \rho \sin(\theta) \sin(\varphi)$$

$$\tan(\theta) = \frac{y}{x}$$

$$z = \rho \cos(\varphi)$$

$$\cos(\varphi) = \frac{z}{\rho}$$

$$dV = \rho^2 \sin(\phi) d\rho d\theta d\phi$$

**Second Derivative Test:** Let  $z = f(x, y)$  be a function of two variables for which the first- and second-order partial derivatives are continuous on some disk containing the point  $(x_0, y_0)$ . Suppose  $f_x(x_0, y_0) = 0$  and  $f_y(x_0, y_0) = 0$ . Define the quantity

$$D = f_{xx}(x_0, y_0)f_{yy}(x_0, y_0) - (f_{xy}(x_0, y_0))^2$$

- i. If  $D > 0$  and  $f_{xx}(x_0, y_0) > 0$ , then  $f$  has a local minimum at  $(x_0, y_0)$ .
- ii. If  $D > 0$  and  $f_{xx}(x_0, y_0) < 0$ , then  $f$  has a local maximum at  $(x_0, y_0)$ .
- iii. If  $D < 0$ , then  $f$  has a saddle point at  $(x_0, y_0)$ .
- iv. If  $D = 0$ , then the test is inconclusive.

**Change of Variables:** We have  $T : S \rightarrow R$  and  $T(u, v) = (g(u, v), h(u, v))$ .

$$\text{Jac}(T) = \begin{vmatrix} \frac{\partial(x, y)}{\partial(u, v)} \end{vmatrix} \text{ and}$$

$$\int \int_S f(x, y) dx dy = \int \int_S f(g(u, v), h(u, v)) |\text{Jac}(T)| du dv.$$

1. (15 points) Find and classify all the critical points of

$$f(x, y) = 2x^3 - 6xy + y^2.$$

2. (20 points) Use Lagrange Multipliers to find the maximum and minimum of the following function

$$f(x, y, z) = x - 2y + 2z$$

subject to the constraint

$$\frac{x^2}{4} + y^2 + z^2 = 12.$$

**3.** (10 points) Calculate the integral

$$\iint_B x e^y dA$$

on  $B = [1, 2] \times [0, 3]$ .

4. Let  $E$  be the region bounded below by  $2x^2 + 2y^2 = z$  and above by the plane  $z = 8$ . Consider

$$\int \int \int_E x^2 + y^2 dV.$$

- a) (8 points) If you were evaluating the given integral over  $E$ , would you integrate in Cartesian, Polar, Cylindrical, or Spherical Coordinates? Explain why you choose the coordinate system you did, and express the bounds for the region  $E$  in that system.

- b) (12 points) Evaluate

$$\int \int \int_E x^2 + y^2 dV.$$

5. Let  $E$  be the region given by  $y \geq 0$  and  $z \geq 0$ , and  $x^2 + y^2 + z^2 \leq 9$ .
- a) (8 points) If you were integrating over  $E$ , would you integrate the region in Cartesian, Polar, Cylindrical, or Spherical Coordinates? Explain why you choose the coordinate system you did, and express the bounds for the region  $E$  in that system.
- b) (12 points) Use integration to find the volume of  $E$ .

6. Answer the following short answer questions.

a) (5 points) A continuous function  $f(x, y)$  will have absolute extrema on a set  $D$  if  $D$  satisfies two conditions. What are those conditions?

b) (5 points) If the average value of  $f(x, y)$  is 5, and  $\int \int_D f(x, y) dA = 30$ , what is the area of a region  $D$ ?

c) (5 points) Find the Jacobian of the transformation  $x(u, v) = u^2 + v$ ,  $y(u, v) = u + v^2$ .