

NAME _____

Rec. Instructor: _____

Signature _____

Rec. Time _____

CALCULUS III - FINAL EXAM

Show all work for full credit. No books or notes are permitted.

Problem	Points	Possible	Problem	Points	Possible
1		9	6		20
2		20	7		15
3		10	8		20
4		20	9		16
5		20	EC		15
Total Score		79			71

Note: Bold letters, like \mathbf{u} , are considered vectors unless specified otherwise.

You are free to use the following formulas on any of the problems.

Projection: $\text{proj}_{\mathbf{u}}\mathbf{v} = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\|^2}\mathbf{u}$

Cylindrical Coordinates:

$$x = r \cos(\theta)$$

$$r = \sqrt{x^2 + y^2}$$

$$y = r \sin(\theta)$$

$$\tan(\theta) = \frac{y}{x}$$

$$z = z$$

$$z = z$$

$$dV = r dr d\theta dz$$

Spherical Coordinates:

$$x = \rho \cos(\theta) \sin(\varphi)$$

$$\rho = \sqrt{x^2 + y^2 + z^2}$$

$$y = \rho \sin(\theta) \sin(\varphi)$$

$$\tan(\theta) = \frac{y}{x}$$

$$z = \rho \cos(\varphi)$$

$$\cos(\varphi) = \frac{z}{\rho}$$

$$dV = \rho^2 \sin(\varphi) d\rho d\theta d\varphi$$

Second Derivative Test: Let $z = f(x, y)$ be a function of two variables for which the first- and second-order partial derivatives are continuous on some disk containing the point (x_0, y_0) . Suppose $f_x(x_0, y_0) = 0$ and $f_y(x_0, y_0) = 0$. Define the quantity

$$D = f_{xx}(x_0, y_0)f_{yy}(x_0, y_0) - (f_{xy}(x_0, y_0))^2$$

- i. If $D > 0$ and $f_{xx}(x_0, y_0) > 0$, then f has a local minimum at (x_0, y_0) .
- ii. If $D > 0$ and $f_{xx}(x_0, y_0) < 0$, then f has a local maximum at (x_0, y_0) .
- iii. If $D < 0$, then f has a saddle point at (x_0, y_0) .
- iv. If $D = 0$, then the test is inconclusive.

Trig Identities: $\cos^2(x) = \frac{1}{2} + \frac{1}{2} \cos(2x)$ $\sin^2(x) = \frac{1}{2} - \frac{1}{2} \cos(2x)$

Line Integrals:

$$\int_C f ds = \int_a^b f(\mathbf{r}(t)) \|\mathbf{r}'(t)\| dt$$

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_a^b \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt$$

Surface Integrals:

$$\int \int_S f dS = \int \int_R f(\mathbf{r}(u, v)) \|\mathbf{t}_u \times \mathbf{t}_v\| du dv$$

$$\int \int_S \mathbf{F} \cdot d\mathbf{S} = \int \int_R \mathbf{F}(\mathbf{r}(u, v)) \cdot (\mathbf{t}_u \times \mathbf{t}_v) du dv$$

Green's Theorem:


$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C P dx + Q dy = \int \int_D (Q_x - P_y) dA$$

$$\int_C \mathbf{F} \cdot \mathbf{N} ds = \int \int_D (P_x + Q_y) dA$$

Stokes' Theorem:

$$\int \int_S \text{curl} \mathbf{F} \cdot d\mathbf{S} = \int_C \mathbf{F} \cdot d\mathbf{r}$$

1. (9 points) For this problem we refer to the following diagram, which is drawn to scale:



The vectors \mathbf{A} , \mathbf{B} , \mathbf{C} , \mathbf{D} , and \mathbf{E} all have length five. All of the angles between the vectors are multiples of 45 degrees. Compute the following explicitly:

- $\mathbf{C} \cdot \mathbf{D}$
- $\|\mathbf{B} \times \mathbf{E}\|$
- $\|\mathbf{A} - \mathbf{C}\|$

- 2.** (20 points) Use Lagrange Multipliers to find the maximum and minimum of the following function

$$f(x, y) = 2x + 6y$$

subject to the constraint

$$3x^2 + y^2 = 36.$$

- 3.** (10 points) Let D be the region given by $[-1, 1] \times [1, e]$. Compute the double integral

$$\int \int_D \frac{x}{y} dA.$$

4. Let E be the region such that $1 \leq x^2 + y^2 \leq 25$, $1 \leq z \leq 3$, and $y \leq 0$.

a) (8 points) If you were evaluating the given integral over E , would you integrate in Cartesian, Polar, Cylindrical, or Spherical Coordinates? Explain why you choose the coordinate system you did, and express the bounds for the region E in that system.

b) (12 points) Use integration to find the volume of E .

5. Let $\mathbf{F}(x, y) = \langle ye^x + 6x - 1, e^x + 2y + 3 \rangle$.

a) (10 points) Determine if \mathbf{F} is conservative. If it is conservative, find a potential function. If it is not conservative, explain why it is not.

b) (10 points) Let C be the oriented curve with parameterization

$$\mathbf{r}(t) = \langle t^2 + t, 2 + t \rangle$$

for $0 \leq t \leq 1$. Evaluate

$$\int_C \mathbf{F} \cdot d\mathbf{r}.$$

State any theorems used.

6. (20 points) Calculate the line integral

$$\oint_C (y^2 - 2xy)dx + (x^3y)dy$$

where C is the rectangle with vertices $(0, 0)$, $(3, 0)$, $(3, 2)$, and $(0, 2)$ oriented counterclockwise. State any theorems used.

7. (15 points) Evaluate the integral

$$\int \int_S 1 dS$$

where S is the portion of $x^2 + y^2 = 9$ between $z = 1$ and $z = 3$.

8. (20 points) Compute the integral

$$\int \int_S \operatorname{curl} \mathbf{F} \cdot d\mathbf{S}$$

where $\mathbf{F}(x, y, z) = x\mathbf{i} + (x^2 + y^2)\mathbf{j} + z^2\mathbf{k}$ and S is the paraboloid $z = 6 - x^2 - y^2$ where $z \geq 2$, oriented outward. State any theorems used.

9. Answer the following short answer questions.

a) (2 points) Suppose that $\nabla f(1,0) = 0$, and the discriminant D of f at $(1,0)$ is positive. Do you need to know any more information to determine if $(1,1)$ is a local maximum?

b) (5 points) Find the Jacobian of the transformation $x(u, v, w) = uvw$, $y(u, v, w) = u + v + w$, and $z(u, v, w) = u - v$.

c) (4 points) If f is a function of x and y , and x and y are each functions of r and s use the chain rule to express $\frac{\partial f}{\partial r}$ and $\frac{\partial f}{\partial s}$.

d) (5 points) Find the equation of the tangent plane to $f(x, y) = yx + x^2y^2$ at point $(2, 3)$.

EC. (15 points) Use the Divergence Theorem to evaluate $\int \int_S \mathbf{F} \cdot d\mathbf{S}$
where

$$\mathbf{F}(x, y, z) = \langle y \sin(\pi x), y^2 z, z + xy \rangle$$

and S is the surface of the box with $-1 \leq x \leq 1$, $2 \leq y \leq 5$, and $0 \leq z \leq 1$. Note that all six sides of the box are included in S .